

DDT and HCH Residue Load in Animal's and Mother's Milk of Lakhimpur Kheeri (Rural areas), Uttar Pradesh-India

Ashok Kumar Maurya*, Khushbu Sharma,** P.E. Joseph**

*CSIR-Central Food & Technological Research Institute, Mysore-India-570020 (Present Address)

** School of Chemical Sciences, Department of Chemistry, St. John's College, Agra- India

Abstract- Organochlorine Pesticides (OCPs) such as DDT and HCH, have stable chemical properties and less biodegradability. As a kind of persistent organic pollutants, they have high risk to the environment and human health. Organochlorine pesticide have been used in Uttar Pradesh in agriculture as a seed dresser, in sanitation, in malaria control programme and in livestock to combat ectoparasites. The pesticides applied drift to areas where cattle graze and plant grow. Because of their chemical stability, they accumulate in the lipid-rich tissue of the body. In the body they circulate throughout all compartments and accumulate in adipose fat. Milk can be used as an evaluation index of environmental contamination by these insecticides, although the main objective of this analysis is to monitor the levels of OCPs in animal's & human's milk of rural areas. The levels of the DDT and HCH residues in animal's & human's milk collected from rural areas having intensive sugarcane, wheat & paddy cultivation in Lakhimpur Kheeri were analysed. All samples contained detectable quantities of DDT, HCH and its metabolites. Total HCH residues were high than those of DDT in all the samples. The total concentration of DDT was found lower than the previous studies carried out in India. These results indicate that the overall DDT levels in Lakhimpur Kheeri region is slightly degrading.

Keywords: DDT, HCH, Milk, Gas chromatography, Mass spectrometry

1. Introduction

India, having agriculture-based economy, is one of largest insecticide consumers in the world. Moreover two-thirds of the pesticides consumed are Class I and II pesticides (WHO) which are highly toxic. No wonder a number of studies from India have reported widespread contamination of various milk, food and water sources to be contaminated with these pesticides. In India, largest pesticide consumption has been in the state of Uttar Pradesh, according to the data of 1995-1996

and 1999-2000, produced by Central Insecticide Board and Registration Committee, India (Srivastava, S et al.2008). Introduced in the 1940s, organochlorine pesticides (OCPs) were widely used in agriculture and pest control until research and public concern regarding the hazards of their use led to government restrictions and bans. Despite restrictions and bans on the use of many organochlorine pesticides in the 1970s and 1980s, they continue to persist in the environment today. OCPs, such as DDT and HCH, have stable chemical properties and less biodegradability. HCH and DDT exhibit broad-spectrum toxicity and residual activities. DDT and HCH are banned in India for agriculture but are still used for controls of vectors in public health. HCH was banned for use in agriculture in 1998 and Lindane was recommended in its place (Mukherjee and Gopal, 2003). These toxicants enter the human body through the food chain and cause serious health problems (John et al.,2001). The presence of OCPs residues in food commodities (Mukherjee and Gopal, 1996), water (Agnihotri et al., 1993), Mother's milk (Nair et al., 1996; Okonkwa et al., 1999; Anoop et al., 2006), dairy milk (John et al., 2001; Mukherjee and Gopal, 1995; Nicholas., 2011), Human blood (Waliszewski et al., 2000) and in skin (Due et al., 1998) have been reported in earlier studies. Epidemiological studies provide evidence that exposures to organochlorine pesticide can produce adverse health effects (Kalpana, 1999; Jiawei et al., 2008).

2. Material and methods

A. Site

Lakhimpur Kheri (28° 27' 0" North, 80° 35' 0" East) is a largest district in terms of area (Total Area = 7680 sqm km) in Uttar Pradesh, India, on the border with Nepal. Sugarcane, wheat, pulses, oilseeds, rice, and potatoes are grown and processed in this district, forming the backbone of the local economy. Some of India's largest sugar mills are in the district. Bajaj sugar mill in Gola Gokarnathand, Bajaj sugar

mill in Palia Kalan and Kishan Sarkari mill in Sampurnanagar are the three largest sugar mills in Asia. The study was conducted in five villages of Lakhimpur Kheeri region namely

Govindnagar, Raninagar, Bhanpur Khajuria, Sumen-nagar and Bheera during 2007 – 2009.

Table 1. The effect of pesticide concentration (ppm) on recovery from spiked buffalo milk

Pesticide	Spiking Level (mg/Kg)	Recovery (%)			Mean recovery % (\pm SD)	Average Recovery % (\pm SD)
		R ₁	R ₂	R ₃		
α -HCH	0.05	87.2	84.0	83.5	84.88 \pm 2.00	86.2 \pm 1.30
	0.02	90.9	83.8	82.8	85.75 \pm 4.44	
	0.10	85.4	88.5	88.5	87.45 \pm 1.78	
	0.05	89.0	85.6	89.6	88.04 \pm 2.15	
β -HCH	0.02	85.7	87.9	83.0	85.50 \pm 2.45	87.6 \pm 1.49
	0.10	89.8	93.2	85.5	89.44 \pm 3.85	
	0.05	85.8	87.4	86.9	86.69 \pm 0.81	
δ -HCH	0.02	95.5	92.4	96.0	94.61 \pm 1.95	93.6 \pm 5.10
	0.10	82.9	86.6	85.8	85.08 \pm 1.94	
	0.05	98.9	92.5	97.9	96.395 \pm 3.44	
γ -HCH	0.02	92.8	95.3	93.7	93.92 \pm 1.26	96.4 \pm 3.87
	0.10	90.0	91.1	90.5	90.53 \pm 0.55	
	0.05	95.3	89.7	91.9	92.27 \pm 2.82	
o,p'-DDT	0.02	94.7	99.0	98.3	97.31 \pm 2.30	92.0 \pm 4.93
	0.10	100.9	98.9	99.9	99.89 \pm 1.0	
	0.05	98.8	99.1	95.2	97.68 \pm 2.17	
p,p'-DDT	0.02	89.9	87.6	89.7	89.06 \pm 1.27	96.0 \pm 1.25
	0.10	88.7	85.2	93.9	89.19 \pm 4.37	
p,p'-DDE	0.05	98.0	96.6	97.3	97.29 \pm 0.0	96.0 \pm 1.25
	0.02	96.4	92.5	95.5	94.78 \pm 2.04	
	0.10	96.9	96.0	94.9	95.92 \pm 1.00	

R₁, R₂ & R₃ are the replicates

B. Sampling

The healthy women samples were collected from donors between age groups of 25 – 35 years and all had normal child deliveries. Normal healthy Cow, Buffalo & Goat were chosen for collection of samples. 50 mL of women milk & 200 mL animal milk were collected in glass bottles by manual suction pump. Milk samples were refrigerated at 4 °C until analyzed.

C. Extraction of the sample

The milk samples were extracted following procedure of Environmental Protection Agency Protocol (EPA1985) and (Ejobi et al., 1996) after some modification. The procedure involved denaturation, solvent extraction, centrifugation, extraction of organic layer and concentration.

Briefly, 5 mL milk samples were mixed with anhydrous sodium sulphate to make a free flowing powder and then transferred into a glass extraction column (150x 20 mm). The dry column was eluted with 100 mL of n-hexane with the first 50 mL allowed to stay in contact with the powder for 10-15 min. The elute was collected in pre-weighed round bottom flask. n-Hexane was evaporated off under a rotary vacuum evaporator. The round bottom flask was then weighed again. The difference in weight is the weight of raw fat extracted. The extracted raw fat was cleaned up in a glass column packed with florisil using eluting solvent of n-hexane & dichloromethane in the ratio of 3:1. The flow rate of elution did not exceed more than 3 mL/min. Then the eluting solvent was passed through anhydrous sodium sulfate and concentrated under rotary vacuum evaporator. Final volume was made up to 1 mL n-hexane and injected 1 μ L in gas liquid chromatograph equipped with an electron capture detector (Ni⁶³).




Figure 1. Standard chromatogram of the DDT and HCH isomers

1, α -HCH 2, γ -HCH 3, β -HCH 4, δ -HCH 5, o,p'-DDT 6, p,p'-DDE 7, p,p'-DDT

3. Instrument

A. GC Instrument

The residues were quantitatively analyzed on Gas Chromatograph-Shimadzu 2010 (Shimadzu, Kyoto, Japan) equipped with split/splitless auto-injector model AOC-20i. The non-polar stationary phase used was a fused silica capillary column DB-5 (5 % phenyl polysiloxane) of 30 m,

0.25 mm i.d and 0.25 μm film thickness (J&W Agilent Palo Alto, CA, USA). GC Solution software was used for instrument control and data analysis.

B. GC/MS Instrument

The residues were further confirmed on Gas Chromatograph-Mass Spectrometer-Quadrupole on electron ionization (EI) mode (Shimadzu 2010, Kyoto, Japan) equipped with with split/splitless auto-injector model AOC-20i. The non-polar stationary phase used was a fused silica capillary column DB-1 (1 % phenyl polysiloxane) of 30 m, 0.25 mm i.d., and 0.25 μm film thickness purchased from J&W Agilent Palo Alto, CA, USA. GCMS Solution software was used for instrument control and data analysis.

4. Analyte recovery and quality control

Milk samples (5 mL) were spiked with the Organochlorine insecticides α -HCH, β -HCH, γ -HCH, δ -HCH, op' DDT, pp'DDD and pp' DDE at 0.02, 0.05, 0.1 $\mu\text{g L}^{-1}$ levels (Table 1). The recovery experiment was performed at the three concentrations and each concentration was analyzed in triplicate. The recovery percentage and standard deviation of organochlorine pesticide are summarized table 3, ranging from 86.2 ± 1.30 to 96.0 ± 1.25 across the three concentrations. The use of milk for recovery studied has earlier has reported by Kanja et al (1983); Ip and Phillips (1986); and Schinas et al. (2000); and Anoop et al. 2006. The limits of detection (LOD) and limit of quantification (LOQ) for OCPs was 0.001 $\mu\text{g/mL}$

Table 2 Residues of DDT contaminants obtained in animal & mother's milk

Milk's type	No of samples	Residues in ppm			
		op'DDT	pp' DDE	PP'DDD	Total
Woman	10	R: (0.053-0.067)	R: (0.046-0.058)	R: (0.052-0.056)	$\Sigma=0.166(\pm 0.002)^*$
		M \pm 0.060 (± 0.009)*	M \pm 0.052 (± 0.010)*	M \pm 0.054 (± 0.007)*	
Cow	10	R: (0.036-0.044)	R: (0.038-0.050)	R: (0.032-0.042)	$\Sigma=0.121(\pm 0.003)^*$
		M \pm 0.040 (± 0.005)*	M \pm 0.044 (± 0.008)*	M \pm 0.037 (± 0.010)*	
Buffalo	10	R: (0.027-0.039)	R: (0.026-0.038)	R: (0.028-0.040)	$\Sigma=0.099(\pm 0.001)^*$
		M \pm 0.033 (± 0.010)*	M \pm 0.032 (± 0.007)*	M \pm 0.034 (± 0.009)*	
Goat	10	R: (0.024-0.032)	R: (0.027-0.039)	R: (0.022-0.034)	$\Sigma=0.089(\pm 0.020)^*$
		M \pm 0.028 (± 0.007)*	M \pm 0.033 (± 0.006)*	M \pm 0.028 (± 0.005)*	

R: range; M= mean; * figures in parenthese gives SD values.

Table 3 Residues HCH contaminants obtained in animal & mother's milk

Milk's type	No of samples	Residues in ppm				Total
		α -HCH	β -HCH	γ -HCH	δ -HCH	
Woman	10	R: (0.045-0.060)	R: (0.37-0.074)	R: (0.032-0.048)	R: (0.021-0.031)	$\Sigma=0.175 (\pm 0.010)^*$
		M \pm 0.053 (± 0.012)*	M \pm 0.056 (± 0.015)*	M \pm 0.040 (± 0.012)*	M \pm 0.026 (± 0.008)*	
Cow	10	R: (0.030-0.050)	R: (0.030-0.046)	R: (0.030-0.040)	R: (0.017-0.027)	$\Sigma=0.135 (\pm 0.005)^*$
		M \pm 0.040 (± 0.016)*	M \pm 0.038 (± 0.012)*	M \pm 0.035(± 0.005)*	M \pm 0.022 (± 0.009)*	
Buffalo	10	R: (0.036-0.046)	R: (0.030-0.042)	R: (0.038-0.052)	R: (0.015-0.025)	$\Sigma=0.142 (\pm 0.009)^*$

		M ±0.041 (±0.006)*	M ±0.036 (±0.008)*	M ±0.045 (±0.012)*	M ±0.020 (±0.008)*	
Goat	10	R: (0.035-0.065) M ±0.050 (±0.014)*	R: (0.025-0.045) M ±0.035 (±0.015)*	R: (0.022-0.036) M ±0.029 (±0.012)*	R: (0.015-0.023) M ±0.019 (±0.007)*	Σ=0.133 (±0.011)*

R: range; M= mean; * figures in parenthese gives SD values.

Table 4 Concentration (ng/g) of organochlorine pesticide residues in human milk in various parts of world

Contries	α-HCH	β-HCH	γ-HCH	δ-HCH	DDE	DDD	DDT	ΣDDT	References
Spain	34.2	235	10.5	279.7	604.1	-	12.5	659.8	Hernandez et al., 1993
Poland	17.5	92.5	15	125	610	12.5	47.5	670	Czaja et al.,1997
Turkey	60	380	17	457	2013	-	100	2357	Coke et al., 1997
France	52	287	37	376	2183	-	79	2262	Bordet et al., 1993
Delhi	1125	495	2100	175	1680	5250	4000	26050	Nair et al., 1996
Mumbai	14.82	259.5	17.5	289.75	232	35	288.5	510.5	Sharma et al.,2001
Agra*									
Bakhoti	32	40	51	123	56	65	58	179	
Chiraigaon	34	43	51	128	56	60	54	170	Anoop et al., 2006
Ghodhakhas	37	40	54	131	56	63	55	174	
Minahas	36	39	52	127	56	66	57	179	
Lakhimpur									
Kheeri	53	56	40	26	60	52	54	166	Present

*Agra:Bakhoti, Chiraigaon, Ghodhakhas & Minahas

and 0.01 µg/mL, respectively. All the solvents & chemical used in the extraction and clean up procedure were special analytical grade for pesticide residues (E.Merk India Ltd.). Pesticide standards were obtained Sigma-Aldrich/ Riedel-de-Haen (Zwijndrecht, The Netherlands).

5. Results and discussion

Total Concentrations of DDTs and HCHs in human breast milk were high than animal milk. This indicates that the residents living in this area have been exposed to relatively high levels of DDTs and HCHs through animal milk, meat, water and agricultural products. o,p'-DDT, p,p'-DDD and p,p'-DDD were analyzed and detected 90% of human's milk, 80% of buffalo's & cow's milk and 70 % of goat's milk. α-HCH, β-HCH, γ-HCH and δ-HCH were also analyzed and found 90% of human milk, 80% of buffalo, cow and goat milk. The concentration of DDT and HCH are presented in Table 2 and 3. Dominant pesticide in all samples examined was HCH in

mother's, buffalo's, cow's, and goat's milk in the concentrations of 175, 135, 142 and 133 ng/ml, respectively. Total DDT levels were found to be 166 ng/ml in mother's milk, 121 ng/ml in cow's & 99 ng/ml buffalo's & 89 ng/ml in goat's milk. These residues show that DDT & HCH used for pest control & agricultural purpose accumulates in human and animal body through the food chain and environment and is excreted through milk. Organochlorine pesticides (OCPs) with their high persistence in the environment accumulate in fatty foods and human adipose tissues. Contamination of human milk by organochlorine and other related compounds has been reported through out the world (GEMS, 1998). During the recent decade, investigations on persistent pollutant (POPs) pollution in the Asian regions and found that relatively high residue levels of DDTs and HCHs exist in food stuffs (Kannan et al., 1997), mussels (Monirith et al., 2003) and avian species (Kunisue et al., 2003) from some developing countries and these contaminants are possibly in use for public health purposes even now. Among Asian developing countries, concentrations of DDTs in human breast milk from Vietnam, mainland China, Cambodia, and Malaysia were

relatively higher than those from other countries (Kunisue et al., 2004). Human milk, at the top of the food chain represents the major route of elimination of OCPs by lactating women (Rogan et al., 1986; Sim and Neil, 1992; IARC, 1991) concluded that there is insufficient evidence in humans but sufficient evidence in experimental animals to classify DDT as a possible carcinogenic to humans. However, body loads of DDT also raise concerns about potential effects on developing infants and children because DDT transfers across the placenta from mother to fetus and exposure continues through breast feeding after birth (Shen et al., 2007). It is well known that they are very dangerous if ingested as an overdose but there is also biological evidence that chronic low-grade exposure to these chemicals, which are very easily absorbed into the body through the skin and lungs, may have adverse effects on mental health. The results obtained from other monitoring studies of organochlorine pesticide in human milk in India and abroad are compared with the results obtained from the present monitoring study in Table 4. α -HCH and β -HCH were found high than previous study of Agra region (Anoop et al. 2006). Presence of high level α -HCH & β -HCH indicate that formulation of these isomers (HCH) was used more frequently in past. Total DDT levels in the present study are less than those reported in India. This indicates gradual phasing out of these compounds in India and has resulted in the reduction of their residues in mother's milk.

6. Conclusion

We have reviewed the available data/information of organochlorine pesticides contamination in rural areas of animal & breast milk from Lakhimpur Kheeri, U.P-India. The result demonstrates that considerable amount of DDT & HCH residues are transferred from the animal & mother to infant. Milk can be considered as a suitable indicator for monitoring the burden of persistent lipophilic chlorinated insecticides in the environment & human body. In view of our observations suggest that further investigation on animal and human exposure in organochlorine pesticides are needed to elucidate future pollution trends and to assess specially infant health risk.

Acknowledgment

Authors are thankful to the Former Principal and Head, Department of Chemistry, St. John's College, Agra (UP), India, for providing research facilities during 2007- 2009.

References

- [1] S.Srivastava, S.S Narvi and S.C Prasad, "Organochlorines and Organophosphates in Bovine Milk Samples in Allahabad Region", International Journal of Environmental Research, vol. 2, No. 2, Spring 2008, pp. 165-168
- [2] I. Mukherjee and M. Gopal, "Insecticide residues in baby food, animal feed and vegetable by gas liquid chromatography," Bull Environ Monit Assess, vol.76, 2003, pp. 185-193.
- [3] P.J. John, N. Bakore and P Bhatnagar, "Assessment of organochlorine pesticide residues levels in dairy milk and buffalo milk from Jaipur city, Rajasthan India," Environ Int, vol. 26, 2001, pp. 231-6.
- [4] I Mukherjee and M.Gopal, "HCH and DDT residues in animal feed and its composition in and around Delhi," Indian Journal Plant Protection, vol. 23, 1996, pp. 61-63.
- [5] N.P Agnihori, V.T Gajbhiye, M Kumar, S.P Mohapatra. Organochlorine insecticide residue in Ganga river water near Farrukhabad, India", Environ Monit Assess, vol 32, 1993, pp.105-112.
- [6] A. Nair, R. Mandapati, P. Dureja, M.K.K. Pillai "DDT and HCH load in mothers and their infants in Delhi, India," Bull Environ Contam Toxicol, vol. 56, 1996, pp.58-64.
- [7] J.O. Okonkwa, L.Kampira, D.D.K. Chinkagule, "Organochlorine insecticide residues in human milk : a study of lactating mother in Sphofaneni, Swziland," Bull Environ Contam Toxicol, vol. 63, 1999, 243-270.

- [8] I Mukherjee and M.Gopal, "HCH and DDT residues in animal feed and its composition in and around Delhi," *Indian Journal Plant Protection*, vol. 23, 1995, pp. 61-63.
- [9] Nicholas, Mwenda Jacob "Assessment of organochlorine pesticide residues in cow milk, sediments and water from Nairobi West region" *Earth Science Frontiers* Volume 15, Issue 5, September 2008, pp- 242-247
- [10] S.M. Waliszewski, A.A. Aruirre, R.M. Infanzon, L. Torres-Sanchez., "Comparison of organochlorine pesticide levels in adipose tissue and blood serum from mother living in Veracruz, Mexico," *Bull Contam Toxicol* , vol. 64, 2000, pp. 8-15.
- [11] B.Kalpana Human health risk assessment for exposures to pesticide: a case study of endocrine disrupters. Proc eeding of the Eighth National Symposium on Environment, Kalpakkam, India",1999, pp. 70.
- [12] S.Gebremichael, T.Birhanu, D.A Tessema OCPs in human a and cow milk and cow's milk in the towns of Asendabo, Serbo and Jimma in South-Western Ethiopia *Chemosphere*, vol. 90/5, February 2013, pp. 1652-1657
- [13] S.P. Mohapatra, M. Kumar, V.T.Gajbhiye, N.P. Agnihotri, "Ground water contamination by organochlorine insecticide residues in a rural area in Indo-Gangetic plains," *Environ Monit Assess*, vol. 35,1995, pp. 155-164.
- [14] Environmental Protection Agency Protocol,1985
- [15] F. Ejobi, L.W Kanja, M.N Kyule, P. Muller, J. Kruger and A.A.R. Latigo. "Organochlorine pesticide residues in mother's milk in Uganda", *Bull Environ.Contam. Toxicol.* vol. 56, 1996, pp. 873-880.
- [16] L.Kanja., J.U.Skaare, I.Nafstad, C.K. Maitai, P. Lokken, "Organochlorine pesticide in human milk from different areas of Kenya" *J Toxicol Environ Helath*, vol. 19, 1983, pp. 449-64
- [17] I. P. HM and J.H. Phillips. "Organochlorine chemical in human breast milk in Madrid Spain, *Arch Environ.Contam. Toxicol*," vol. 150, 1993, pp.308
- [18] V. Schinas, M. Leotsinidis, V. Tsapanos, X.G. Kondakis, "Organochlorine pesticide residues in human body milk from south west Greece : association weakly food consumption pattern of mother," *Arch Environmental Health* 2000; vol. 556, pp. 411-7.
- [19] A. Kumar, P. Dayal , G. Shukla, G. Singh and P.E. Joseph " DDT and HCH residue load in mother's breast milk: A survey of lactating mother's from remote villages in Agra region." *Environment International*, vol 32 , 2006, pp.248-251
- [20] G.E.M.S, Food Global Environment Monitoring System-Food Contamination Monitoring Organization (Press Release WHO),vol.10, 1998..
- [21] K.Kannan,, S.Tanabe, J.P.Giesy, R.Tatsukaw, "Organochlo -rine pesticides and polychlorinated biphenyls in foodstuffs from Asianand Oceanic countries," *Reviews of Environmental Contamination & Toxicology*, vol.152,1997, pp. 1-55
- [22] I. Monirith, D. Ueno, S. Takahashi, H Nakata, A. Sudaryanto, A. Subramanian, S Karuppiah, Ismail,, Muchtar, M., Zheng, J.,Richardson, B.J., A.Prudente, N.D Hue, T.S Tana, A.V. Tkalin, S.Tanabe, "Asia-Pacific mussel watch: monitoring contamination of persistent organochlorinecompounds in coastal waters of Asiancountries, *Marine Pollution*" *Bulletin*, vol. 46, 2003, pp.281-300.
- [23] T. Kunisue, M. Watanabe, A. Subram, "Accumulation features of persistent

- organochlorines in resident and migratory birds from Asia”, Environmental pollution,” vol.125, 2003, pp. 157-172.
- [24] T.Kunisue, M.Someya, M.Watanabe,T.Toyota, Y Kuroda, J. Nagayama, S. Tanab, (“Contamination by persistent organic pollutants (POPs) in human breast milk collected from primiparae in Ehime prefecture”, Environmental Science ,vol. 17, 2004, pp 37-48.
- [25] W.J. Rogan, B.C. Gladen, J.D. McKinney, N. Carreras, P. Hardy, J. Thullen, J. Tinglestad, M. Tully, “Neonatal effects of transplacental exposure to PCBs and DDE, Journal of Pediatrics,” vol.109, 1986 , pp. 335-341.
- [26] M.R. Sim, J.J. Mc Neil, “Monitoringchemical exposures using breast milk: amethodological review,” American Journal of Epidemiology, vol. 136, 1992, pp.1-11.45.
- [27] I.A.R.C, International Agency for Researchon Cancer, “DDT and Associated Compounds,Summaries and Evaluation1”,vol.7,1991.
- [28] S.Srivastava, S.S Narvi and S.C Prasad, “ Organochlorines and Organophosphates in Bovine Milk Samples in Allahabad Region International Journal of Environmental Research”, Vol. 2, No. 2, Spring 2008, pp. 165-168
- [29] H. Shen, K.M. Main, H.E. Virtanen, I.N. Damggard, A.M. Haavisto, M.Kaleva, K.A.Boisen, I.M Schmidt, M. Chellakooty, N.E. Skakkebaek, J. Toppari, K.W Schramm, “From mother to child: investigation of prenatal and postnatal exposure to persistent bioaccumulating toxicantsusing breast milk and placenta biomonitoring,” Chemosphere, vol. 67, 2007, pp. 256-262.